

Selected Topics

1. Vibrational coupling
2. Electronic transport

Theoretical investigation of local vibrational excitations by STM-IETS in molecules

***Jessica Walkenhorst (1); Marius Wanko (1); Alberto Castro (2); Angel Rubio (1)**

1. Nano-Bio Spectroscopy group and ETSF Scientific Development Centre, Dpto. Física de Materiales, Universidad del País Vasco, Centro de Física de Materiales CSIC-UPV/EHU-MPC and DIPC - San Sebastián, Spain; 2. Institute for Biocomputation and Physics of Complex Systems (BIFI), University of Zaragoza, Spain

Corresponding Author: Jessica Walkenhorst (walkenho@gmail.com)

Tunneling electrons can be used to excite vibrations in molecules. By combining the high spatial resolution of scanning electron microscopy (STM) with the chemical selectivity of Inelastic Electron Tunneling Spectroscopy (IETS) it is now possible to obtain vibrational spectra of single molecules adsorbed on a surface. This way identification and selective removal of isotopes in molecules can be done (e.g. H and D in acetylene).

Interestingly in large molecules, the vibrational spectrum obtained by STM-IETS seems to depend on the impact point of the electrons on the molecular surface.

Selective excitation of vibrational modes by the STM tip has been proposed to explain these experimental findings (so-called 'excitation of local vibrations' model [1]). Therefore, we want to simulate the dynamical response of acetylene and phtalocyanine (NPc) induced by the impact of tunneling electrons. To this end, we perform molecular dynamics simulations based on time-dependent density functional theory (TDDFT) in which tunneling electrons are modeled by means of Gaussian wave packets. Vibrational spectra are then obtained by Fourier analysis of the velocity autocorrelation function. This way, we study the relative strength of the excited molecular vibrations in dependence on the impact point of the tunneling electrons. Finally, a comparison between numerical results and experimental findings is attempted and the validity of the 'excitation of local vibrations' model is discussed.

This work was supported by the Spanish MEC (FIS2007-65702-C02-01), ACI-PPomociona (ACI2009-1036), Grupos Consolidados UPV/EHU del Gobierno Vasco (IT-319-07), ETORTEK (inanoGUNE), the European Union through e-I3 ETSF project (Contract Number 211956). We acknowledge support by the Barcelona Supercomputing Center, Red Espanola de Supercomputacion.

[1] N. Ogawa, G. Mikaelian and W. Ho, Phys. Rev. Lett. **98**, 166103 (2007)