

## Contribution submission to the conference Dresden 2014

### **Correlated Light-Matter Interactions in Cavity QED —**

•JOHANNES FLICK<sup>1</sup>, HEIKO APPEL<sup>1</sup>, and ANGEL RUBIO<sup>1,2</sup> — <sup>1</sup>Fritz-Haber-Institut der Max-Planck-Gesellschaft, Berlin, Germany — <sup>2</sup>NanoBio Spectroscopy group and ETSF, Universidad del País Vasco, San Sebastián, Spain

In the last decade, time-dependent density functional theory (TDDFT) has been successfully applied to a large variety of problems, such as calculations of absorption spectra, excitation energies, or dynamics in strong laser fields. Recently, we have generalized TDDFT to describe electron-photon systems (QED-TDDFT) [1, 2]. Here, matter and light are treated on an equal quantized footing.

In this work, we present the first numerical calculations in the framework of QED-TDDFT. We present exact solutions for fully quantized prototype systems consisting of atoms or molecules placed in optical high-Q cavities and coupled to quantized electromagnetic modes. We focus on the electron-photon xc-contribution by calculating exact Kohn-Sham potentials using fixed-point inversions and present the performance of the first approximated xc-potential based on an optimized effective potential (OEP) approach.

[1] I. Tolkatly, Phys. Rev. Lett. **110**, 233001 (2013).

[2] M. Ruggenthaler et.al., Phys. Rev. A **90**, 012508 (2014).

**Part:** O  
**Type:** Vortrag; Talk  
**Topic:** Focussed Session: Frontiers of Electronic Structure Theory  
**Email:** flick@fhi-berlin.mpg.de